

Where the Links–Gould invariant first fails to distinguish nonmutant prime knots

David De Wit and Jon Links*

February 1, 2008

Abstract

It is known that the first two-variable Links–Gould quantum link invariant $LG \equiv LG^{2,1}$ is more powerful than the HOMFLYPT and Kauffman polynomials, in that it distinguishes all prime knots (including reflections) of up to 10 crossings. Here we report investigations which greatly expand the set of evaluations of LG for prime knots. Through them, we show that the invariant is complete, modulo mutation, for all prime knots (including reflections) of up to 11 crossings, but fails to distinguish some nonmutant pairs of 12-crossing prime knots. As a byproduct, we classify the mutants within the prime knots of 11 and 12 crossings. In parallel, we learn that LG distinguishes the chirality of all chiral prime knots of at most 12 crossings. We then demonstrate that every mutation-insensitive link invariant fails to distinguish the chirality of a number of 14-crossing prime knots. This provides 14-crossing examples of chiral prime knots whose chirality is undistinguished by LG .

1 The Links–Gould invariant

For any positive integers m and n , the Links–Gould quantum link invariant $LG^{m,n}$ is a two-variable invariant of oriented links. We here describe it using the variables q and $p \equiv q^{\alpha+(m-n)/2}$ which are inherited from its definition via the 2^{mn} -dimensional α -parametric representation of highest weight $(\dot{0}_m \mid \dot{\alpha}_n)$ of the quantum superalgebra $U_q[gl(m|n)]$. Its construction was originally described in [12, 25], and some of its properties together with some concrete evaluations are available in [5, 6, 8, 9].

The case $LG^{1,1}$, in which the two variables degenerate to a single variable (p), is actually the Alexander–Conway polynomial, and the next simplest case, $LG^{2,1}$, is the first truly two-variable invariant of the family. Apart from $LG^{1,1}$, more is known about $LG^{2,1}$ than about any of its kin [4, 15–21], and we generally refer to $LG^{2,1}$ as *the* Links–Gould invariant LG . In particular, it has been proven to be polynomial [17]; this is still only surmised for general $LG^{m,n}$.

*Department of Mathematics, The University of Queensland, 4072, Brisbane, Australia.
Dr.David.De.Wit@yahoo.com.au, jrl@maths.uq.edu.au

As with all such quantum link invariants, $LG^{m,n}$ is insensitive to the inversion of links, and this manifests in its invariance under the inversion of p (modulo a sign — this is a well-known property of the Alexander–Conway polynomial). That is, for any oriented link L , the (polynomial) evaluation $LG_L^{m,n}$ is necessarily palindromic in p , meaning that $LG_{-L}^{m,n}(q, p) \doteq LG_L^{m,n}(q, p^{-1})$, where by $-L$ we intend the inversion of L , and by $\doteq$ we intend equality up to a sign ± 1 . Also, as for most of the more familiar link invariants, $LG^{m,n}$ fails to distinguish mutant links.

As we know these limitations, it is of interest to determine where LG first fails to distinguish nonmutant prime knots. Significantly, LG is known to be complete for all prime knots (including reflections) of up to 10 crossings [4]. In this, it is known to be more powerful than the HOMFLYPT and Kauffman polynomials. Unlike the situation for those polynomials, however, a set of skein relations sufficient to determine LG for any arbitrary link is not yet known, and most evaluations to date have been performed using a computationally expensive state model method.

Here, we report on evaluations of LG for a range of the prime knots from the Hoste–Thistlethwaite–Weeks (HTW) tables of prime knots of up to 16 crossings [13]. The present list of evaluations expands the previous published list [4] by a factor of 200. Of significance, we determine that LG is complete, modulo mutation, for all prime knots (including reflections) of up to 11 crossings, but that it fails to distinguish several nonmutant 12-crossing prime knots.

A byproduct of these investigations is a listing of all mutant cliques within the prime knots of 11 and 12 crossings, which should serve as a useful test-bed for future experiments with invariants that are not mutation-insensitive. We also mention a number of open basic questions about mutation.

It is also known that LG is not always able to distinguish between the reflections of chiral links, as Ishii and Kanenobu have constructed an infinite family of chiral knots whose chirality is undetected by LG [20]. At least some of these links are prime knots, although of unknown crossing numbers. Herein, we demonstrate that LG distinguishes the chirality of all chiral prime knots of up to 12 crossings. We also identify several 14-crossing chiral prime knots whose chirality is not distinguished by LG , or indeed by any mutation-insensitive invariant. That said, we have as yet no example of a chiral link whose chirality is not detected by LG yet is detected by either the HOMFLYPT or the Kauffman polynomial.

2 Evaluations of LG for the HTW prime knots

Our evaluations of LG are with respect to the HTW tables of prime knots of up to 16 crossings [13]. Dowker–Thistlethwaite (DT) codes and many other data associated with the knots in these tables may be accessed via the program KNOTSCAPE (version 1.01), as mentioned in [13]. We let c_i^A (respectively c_i^N) denote the i th alternating (respectively nonalternating) prime knot (type) of c crossings in the HTW tables. The presence of the superscript decorating the

crossing number identifies the knot as from the HTW tables, and facilitates the concurrent use of the classical undecorated Alexander–Briggs notation; for instance 5_1^A and 5_2 denote the same knot type. In [4] is found a listing of evaluations of LG for all prime knots of up to 10 crossings, listed with respect to the Alexander–Briggs ordering. Herein, we abandon that ordering, and instead adopt the HTW ordering.

In the classical tables, as reproduced for instance in [24], canonical representatives of knot types are described graphically, however these diagrams omit an orientation, so only describe *unoriented* knots. In contrast, the HTW tables describe knot types via DT codes, and as any DT code necessarily encodes both a link and its reflection, the tables do not prescribe chiralities (orientations of space), although they do prescribe string orientations. (As quantum link invariants are insensitive to inversion, this is unimportant to our considerations.) An evaluation of a link invariant for a link described in terms of a DT code thus should be regarded as unique up to a symmetry transformation corresponding to that of reflection of the link. For $LG^{m,n}$ this transformation corresponds to the inversion of the variable q . Using the HTW tables to define knots thus means that LG polynomials should be regarded as attached to knot *types*, rather than knots per se. Herein, the statement that LG cannot distinguish distinct links L_1 and L_2 means that the polynomials LG_{L_1} and LG_{L_2} are either equal or related by inversion of q . In this situation, we describe the links as *LG-equivalent*. Thus the LG -polynomial of an achiral link is necessarily palindromic in q .

Our evaluation of some $LG^{m,n}$ for a particular link L is generally obtained by the application of a state model method to a braid β , where the closure $\widehat{\beta}$ corresponds to L . We have implemented this in a MATHEMATICA package called LINKS–GOULD EXPLORER; a description of the algorithms used appears in [3]. Evaluations of $LG^{2,1}$ are currently generally feasible for braids of string index at most 5 (*sometimes* 6 and even 7) and a ‘reasonable’ number of crossings (say less than 30). By ‘feasible’, we intend computations which demand at most about 1Gb of memory and run-times of at most a few CPU-hours on current commonly-available hardware. Memory requirements increase explosively with braid width.

Braids for the HTW knots are obtainable via the implementation of Vogel’s Algorithm [29] in the MATHEMATICA package K2K (version 1.3.3) by Imafuji and Ochiai [14]. (K2K is an interface to an underlying C program **KnotTheoryByComputer** by Ochiai.) Unfortunately, a bug in the implementation of Vogel’s Algorithm in K2K means that it fails to terminate for exactly 11 of the 1,701,936 HTW knots. (Otherwise, it yields *correct* braids!) In [2] we explain the application of other methods to determine braids for these 11 knots; the net result is that we do have braids for all the HTW knots.

Recall that a *minimal* braid is one whose width is minimal over all candidate braids for a given knot, and this width is called the *string index* of the knot. Sadly, the automatically-generated K2K-braids are generally far from minimal, and where possible, we have used thinner ones. Specifically, we have to hand minimal braids for all the HTW knots of up to 12 crossings. This data is obtained from Livingston’s website TABLE OF KNOT INVARIANTS [26] together

with that of Stoimenow, called KNOT DATA TABLES [28]. The braids for knots of up to 10 crossings were constructed by Gittings [10] and those of 11 and 12 crossings by Stoimenow. These braids are also (generally) of minimal length for their minimal width, and of a structure which facilitates their use in state model algorithms. For the knots of 11 (respectively 12) crossings, they are of maximum string (that is, braid) index 6 (respectively 7).

We refer to our full collection of braids from these diverse sources as the K2K'-braids. To describe them, we introduce some convenient notation. Where P is either A or N , denote by $\mathbb{K}_c^P$ the set of all prime knots c_i^P of c crossings, and set $\mathbb{K}_c \triangleq \mathbb{K}_c^A \cup \mathbb{K}_c^N$. Similarly, denote by $\mathbb{K}_{c,s}^P$ the set of all prime knots c_i^P of c crossings whose K2K'-braids have s strings, and set $\mathbb{K}_{c,s} \triangleq \mathbb{K}_{c,s}^A \cup \mathbb{K}_{c,s}^N$.

Table 1 lists the sizes of the $\mathbb{K}_{c,s}^P$. An artefact of the algorithm used to construct the K2K-braids is that a K2K-braid corresponding to an HTW prime knot of an odd (respectively even) number of crossings has an even (respectively odd) string index.

We have evaluated LG for a total of 53,418 of these knot types (not counting reflections), including all complete sets $\mathbb{K}_{c,s}^P$ lying above the zigzags in Table 1.

- The state model method has been applied to the K2K'-braids of a total of 48,399 prime knots, namely those from:
 - all $\mathbb{K}_c$ for $c \leq 12$, and
 - all $\mathbb{K}_{c,s}$ for $c \leq 15$ and $s \leq 5$, and $\mathbb{K}_{16,3}$.¹
- A formula of Ishii [16] for LG for 2-bridge knots has been applied to the 5546 (necessarily alternating) 2-bridge knots within the HTW tables (as determined in [7]).

These classes have an overlap of only 527 knots as the K2K'-braids for the 2-bridge knots are generally on more than 5 strings. Thus, we have evaluations of LG for a net $48,399 + 5546 - 527 = 53,418$ prime knots. Table 2 provides a breakdown of these totals with crossing numbers.

For comparison with Table 1, Table 3 presents the numbers of c -crossing 2-bridge knots with s -string K2K'-braids.

We illustrate an LG -polynomial using 15_{139717}^N , the only achiral knot of odd crossing number in the HTW tables, depicted in Figure 1.² Although its K2K-braid is on 6 strings, it is still LG -feasible. We obtain the following polynomial, where we have written $\bar{q} \triangleq q^{-1}$ and $\bar{p} \triangleq p^{-1}$. The polynomial is palindromic in p , as are all LG -polynomials. It is also palindromic in q , as the knot is achiral.

¹The LG -feasible $\mathbb{K}_{16,5}$ is omitted due to its size, as insufficient CPU-years were available.

²This knot is hyperbolic, and has symmetry group D_1 (see §3 and §5).

		C															
		0	3	4	5	6	7	8	9	10	11	12	13	14	15	16	totals
S	1	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1
	2	.	1	.	1	.	1	.	1	.	1	.	1	.	1	.	7
	3	.	.	1	1	2	2	8	4	23	6	71	.	225	.	746	1089
	4	.	.	.	.	1	4	7	23	48	149	290	1017	.	8018	.	9557
	5	.	.	.	.	.	.	3	13	47	164	652	.	6905	.	80342	88126
	6	.	.	.	.	.	.	.	.	5	47	258	2570	.	40388	.	43268
	7	.	.	.	.	.	.	.	.	.	.	17	.	9479	.	190642	200138
	8	.	.	.	.	.	.	.	.	.	.	.	1145	.	30684	.	31829
	9	.	.	.	.	.	.	.	.	.	.	.	.	2728	.	94924	97652
	10	.	.	.	.	.	.	.	.	.	.	.	140	.	5838	.	5978
	11	.	.	.	.	.	.	.	.	.	.	.	.	196	.	12697	12893
	12	.	.	.	.	.	.	.	.	.	.	.	5	.	334	.	339
	13	.	.	.	.	.	.	.	.	.	.	.	.	3	.	448	451
totals		1	1	1	2	3	7	18	41	123	367	1288	4878	19536	85263	379799	491328

Alternating

		C																
		0	3	4	5	6	7	8	9	10	11	12	13	14	15	16	totals	
S	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	0	
	2	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	0	
	3	.	.	.	.	.	.	3	.	14	.	50	.	121	.	478	666	
	4	.	.	.	.	.	.	.	8	26	130	333	1247	.	15559	.	17303	
	5	.	.	.	.	.	.	.	.	2	55	486	.	11103	.	244352	255998	
	6	.	.	.	.	.	.	.	.	.	.	19	3067	.	92169	.	95255	
	7	.	.	.	.	.	.	.	.	.	.	.	.	13589	.	546065	559654	
	8	.	.	.	.	.	.	.	.	.	.	.	768	.	55215	.	55983	
	9	.	.	.	.	.	.	.	.	.	.	.	.	2578	.	203041	205639	
	10	.	.	.	.	.	.	.	.	.	.	.	.	28	.	5021	5049	
	11	.	.	.	.	.	.	.	.	.	.	.	.	.	45	.	14856	14901
	12	.	.	.	.	.	.	.	.	.	.	.	.	.	.	66	.	66
	13	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	114	114
totals		0	0	0	0	0	0	3	8	42	185	888	5110	27436	168030	1008906	1210608	

Nonalternating

		C																
		0	3	4	5	6	7	8	9	10	11	12	13	14	15	16	totals	
S	1	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	
	2	.	1	.	1	.	1	.	1	.	1	.	1	.	1	.	7	
	3	.	.	1	1	2	2	11	4	37	6	121	.	346	.	1224	1755	
	4	.	.	.	.	1	4	7	31	74	279	623	2264	.	23577	.	26860	
	5	.	.	.	.	.	.	3	13	49	219	1138	.	18008	.	324694	344124	
	6	.	.	.	.	.	.	.	.	5	47	277	5637	.	132557	.	138523	
	7	.	.	.	.	.	.	.	.	.	.	17	.	23068	.	736707	759792	
	8	.	.	.	.	.	.	.	.	.	.	.	1913	.	85899	.	87812	
	9	.	.	.	.	.	.	.	.	.	.	.	.	5306	.	297965	303271	
	10	.	.	.	.	.	.	.	.	.	.	.	168	.	10859	.	11027	
	11	.	.	.	.	.	.	.	.	.	.	.	.	241	.	27553	27794	
	12	.	.	.	.	.	.	.	.	.	.	.	5	.	400	.	405	
	13	.	.	.	.	.	.	.	.	.	.	.	.	3	.	562	565	
totals		1	1	1	2	3	7	21	49	165	552	2176	9988	46972	253293	1388705	1701936	

All

Table 1: The sizes of $\mathbb{K}_{c,s}^A$, $\mathbb{K}_{c,s}^N$ and $\mathbb{K}_{c,s}$. Above the zigzags lie sets for which LG has been evaluated using a state model method.

	c																
	0	3	4	5	6	7	8	9	10	11	12	13	14	15	16		totals
state model	1	1	1	2	3	7	21	49	165	552	2176	2265	18354	23578	1224		48399
2-bridge	.	1	1	2	3	7	12	24	45	91	176	352	693	1387	2752		5546
overlap	.	1	1	2	3	7	12	24	45	91	176	26	95	37	7		527
net	1	1	1	2	3	7	21	49	165	552	2176	2591	18952	24928	3969		53418

Table 2: Numbers of c -crossing prime knots for which LG has been evaluated.

		C															
		3	4	5	6	7	8	9	10	11	12	13	14	15	16	totals	
S	2	1	.	1	.	1	.	1	.	1	.	1	.	1	.	7	
	3	.	1	1	2	2	3	3	4	4	5	.	6	.	7	38	
	4	.	.	.	1	4	6	12	15	24	28	25	.	36	.	151	
	5	.	.	.	.	.	3	8	22	40	73	.	89	.	151	386	
	6	.	.	.	.	.	.	.	4	22	60	123	.	276	.	485	
	7	.	.	.	.	.	.	.	.	.	10	.	280	.	736	1026	
	8	.	.	.	.	.	.	.	.	.	.	151	.	592	.	743	
	9	.	.	.	.	.	.	.	.	.	.	.	251	.	1145	1396	
	10	.	.	.	.	.	.	.	.	.	.	.	49	.	400	.	449
	11	.	.	.	.	.	.	.	.	.	.	.	.	64	.	610	674
	12	.	.	.	.	.	.	.	.	.	.	.	3	.	82	.	85
	13	.	.	.	.	.	.	.	.	.	.	.	.	3	.	103	106
	totals		1	1	2	3	7	12	24	45	91	176	352	693	1387	2752	5546

Table 3: The numbers of c -crossing 2-bridge knots with s -string $K2K'$ -braids.

$$\begin{aligned}
& (2\overline{q}^6 - 10\overline{q}^4 + 72\overline{q}^2 + 181 + 72q^2 - 10q^4 + 2q^6) \\
- & (\overline{p}^2 + p^2) (\overline{q}^7 - 4\overline{q}^5 + 8\overline{q}^3 + 118\overline{q} + 118q + 8q^3 - 4q^5 + q^7) \\
+ & (\overline{p}^4 + p^4) (-4\overline{q}^4 + 32\overline{q}^2 + 80 + 32q^2 - 4q^4) \\
- & (\overline{p}^6 + p^6) (7\overline{q}^3 + 30\overline{q} + 30q + 7q^3) \\
+ & (\overline{p}^8 + p^8) (2\overline{q}^4 + 11\overline{q}^2 + 20 + 11q^2 + 2q^4) \\
- & (\overline{p}^{10} + p^{10}) (-\overline{q}^5 + \overline{q}^3 + 10\overline{q} + 10q + q^3 - q^5) \\
+ & (\overline{p}^{12} + p^{12}) (-2\overline{q}^4 + \overline{q}^2 + 7 + q^2 - 2q^4) \\
- & (\overline{p}^{14} + p^{14}) (-\overline{q}^3 + 2\overline{q} + 2q - q^3) \\
+ & (\overline{p}^{16} + p^{16}) (1)
\end{aligned}$$

Note that LG -polynomials often have a regular sign pattern, in that the coefficients within the q -polynomial corresponding to any given power of p are all of the same sign, and this sign alternates with rising (even) powers of p . Examples of this pattern are seen in LG -polynomials presented in §4. It is not known what property of a link causes this pattern. Observe that $LG_{15N_{139717}}$ is a (slight) exception to this rule. (The prime knots of up to 10 crossings with such exceptional LG -polynomials are identified in [4].) It does, however, satisfy another general (and still unexplained) property of LG -polynomials: Each of its q -polynomials contains either only odd or only even powers of q .

We conclude this section with a few remarks about the computability of $LG^{3,1}$, the next available Links–Gould invariant. The $LG^{3,1}$ -polynomials are generally of higher degree than those of $LG^{2,1}$, and are correspondingly more expensive to evaluate. Using our state model method we have only been able to evaluate $LG^{3,1}$ for prime knots in the following complete sets $\mathbb{K}_{c,s}$, which together contain only 380 knot types:




Figure 1: 15^N_{139717} : The first achiral knot of odd crossing number.

- all $\mathbb{K}_c$ for $c \leq 10$, and
- all $\mathbb{K}_{c,s}$ for $c \leq 12$ and $s \leq 3$, and
- $\mathbb{K}_{13,2} \equiv \{13^A_{4878}\}$ and $\mathbb{K}_{15,2} \equiv \{15^A_{85263}\}$.

We find that $LG^{3,1}$ is complete for these knots (including reflections).

3 Mutant cliques within the HTW tables

Recall that a (Conway) mutation of a $(2,2)$ -tangle (and thence of a link in which the tangle is embedded) amounts to a rotation of π about one of three orthogonal axes of the tangle. More generally, the result of a sequence of mutations is regarded as a mutation in itself, and as mutation is reflexive, symmetric and transitive, we may discuss equivalence classes of knots under mutation.

Many link invariants fail to distinguish mutants: the list includes each $LG^{m,n}$ together with all the well-known polynomial link invariants (see §4), and sundry others including the hyperbolic volume (see below) and Khovanov homology [30]. With a view to identifying where LG fails to identify nonmutant prime knots, we here describe a partial classification of the mutants within the HTW tables. Specifically, we classify those within $\mathbb{K}_{11}$ and $\mathbb{K}_{12}$. Of course, we need only describe the equivalence classes of size greater than 1, and these we refer to as *mutant cliques* (within the HTW tables). Although the identification turns out to be relatively straightforward to obtain, this information apparently does not appear in the existing literature. This situation reflects a remarkable lack of research into mutation, the notable exception being the search for link invariants which are sensitive to it, for example [27].

We determine mutant cliques by first using mutation-insensitive invariants to filter the HTW knots into candidate mutant cliques, and then inspect the diagrams to determine if mutations are visible between their elements. The inspection process is nonalgorithmic, and may fail to identify some genuine mutants, and cannot demonstrate when candidates are *not* mutants. Nevertheless, in this manner, we obtain a complete classification of the mutant cliques

within $\mathbb{K}_{11}$ and $\mathbb{K}_{12}$. Moreover, the filtering allows us to make some qualitative statements about the remaining undetermined mutant cliques of higher crossing numbers.

The main invariant we use to determine candidate mutant cliques is the hyperbolic volume, so we digress to review it. To that end, if the complement $S^3 - L$ of a link L can be assigned a hyperbolic metric, it can be decomposed with respect to that metric, and L is called *hyperbolic*.³ In such a case, the complement can be assigned a *hyperbolic volume*, which is an (algebraic) positive real number. This quantity may be regarded as a property of L itself, and is an invariant of L , for the following reason.

The Gordon–Luecke theorem [11] tells us that topologically equivalent (true) knots have homeomorphic knot complements and vice-versa. (Unfortunately, this statement does not hold more generally for multicomponent links.) It follows that the hyperbolic decomposition of the knot complement distinguishes all hyperbolic knots. In particular, the hyperbolic decomposition distinguishes *mutant* hyperbolic knots. Unfortunately however, the hyperbolic *volume* alone does *not* distinguish mutant hyperbolic knots. As composite knots are nonhyperbolic, a mutation of a hyperbolic prime knot is necessarily a (hyperbolic) prime knot, rather than a composite knot. It appears that we do not know whether a mutation of a nonhyperbolic prime knot is a nonhyperbolic prime knot (as opposed to a composite knot). The software available for determining hyperbolic decompositions, and thus hyperbolic volumes, is the SNAPPEA program by Weeks [13]; we use it via its embedding within KNOTSCAPE.

The hyperbolic volume is a powerful invariant, and generally distinguishes many more knots than polynomial invariants, however, distinct knots with the same hyperbolic volume are not *necessarily* mutants. Indeed, it sometimes fails to distinguish between prime knots of different crossing number, for example $(5_1^A, 12_{242}^N)$, which are distinguished by both the HOMFLYPT and the Kauffman polynomials. Moreover SNAPPEA only determines a hyperbolic volume as a decimal approximation (to 10 digits of accuracy and precision) to its true value. Thus, as for polynomial invariants, whilst it can be used to prove that given knots are not mutants, it can only be used as an *indicator* of possible mutants.

³Nonhyperbolic knots are rare in the HTW tables. Including the unknot, there are only 33 of them (out of 1,701,936). They include 13 torus knots, which always form only a small proportion of all prime knots of any given crossing number, and 20 satellite knots (it appears that these will also always be a small number of the total of any given crossing number). They are:

- 13 torus knots, which include the unknot 0_1^A (which may be regarded as the $(2, 1)$ torus knot); the closed odd 2-braids $3_1^A, 5_2^A, 7_7^A, 11_{367}^A, 13_{4878}^A$, and 15_{85263}^A ; and sundries $8_3^N, 9_{41}^A, 10_{21}^N, 14_{21881}^N, 15_{41185}^N, 16_{783154}^N$; together with
- 20 satellites of the trefoil: $13_{4587}^N, 13_{4639}^N, 14_{22180}^N, 14_{26039}^N, 15_{40211}^N, 15_{59184}^N, 15_{115646}^N, 15_{124802}^N, 15_{142188}^N, 15_{156076}^N, 16_{253502}^N, 16_{400459}^N, 16_{696530}^N, 16_{697612}^N, 16_{703714}^N, 16_{703716}^N, 16_{800356}^N, 16_{800378}^N, 16_{958969}^N, 16_{958982}^N$.

Importantly, note that there are no 12-crossing nonhyperbolic knots, and that the only 11-crossing example is easily identified.

We continue the main thread of our discussion with some other salient facts about mutation.

- A mutation of a true knot is a true knot, rather than a multicomponent link. No mutation of a nontrivial knot can create the unknot [1, pp50–51].
- Although a mutation of an alternating projection of an (alternating) knot is always an alternating knot [1, pp50–51], it is apparently not known whether a mutation of a nonalternating projection of an alternating knot is always an alternating knot. However, at least for prime knots of low crossing numbers, mutant cliques appear to only contain knots of common ‘alternatingness’.
- Mutation certainly preserves the numbers of crossings of link *projections*, and it can of course leave a link unchanged. It appears that the question of whether mutation can convert a chiral link into its reflection is still unanswered; in any case we know no explicit examples of this phenomenon, and as we are only considering knot *types*, we may ignore this issue.

The first point at which mutation changes a prime knot into a distinct knot is within $\mathbb{K}_{11}$ [24, p44] (recall the Kinoshita–Terasaka pair $(11_{34}^N, 11_{42}^N)$). In fact, at least for prime knots of low crossing numbers, mutation often preserves crossing number. This may not more generally be the case, as crossing number is not an invariant defined by a symmetry.

To determine candidate mutant cliques within the HTW tables, we first reduce the full set of HTW knots into equivalence classes of common (apparent) hyperbolic volume, and then further reduce these equivalence classes with the HOMFLYPT and Kauffman polynomials, only retaining sets of size greater than 1. (For good measure, we further filter the candidate sets with LG where possible.) By inspection, each such candidate mutant clique lies within a particular class $\mathbb{K}_c^P$, so every genuine mutant clique also lies within a class $\mathbb{K}_c^P$. For the prime knots of up to 16 crossings, mutation preserves crossing number and alternatingness.

In this manner, within $\mathbb{K}_{11}$ we determine 16 mutant candidate pairs, all of which actually are mutants. These are depicted in Figures 2 and 3. In many of the pictures, a mutation can be seen by immediate inspection; others require a little more work. We mention that we describe the mutant cliques of knot types, ignoring reflections, so the identification of a mutation relating pairs of diagrams may include imposing a reflection on one of them. The inspection process is in some cases made easier by the fact that nonalternating mutant cliques sometimes share the same shadows (crossing-information-obliterated diagrams) as alternating mutant cliques — compare $(11_{24}^A, 11_{26}^A)$ with $(11_{39}^N, 11_{45}^N)$, $(11_{40}^N, 11_{46}^N)$ and $(11_{41}^N, 11_{47}^N)$ — and this in turn reflects the way in which the HTW tables are compiled.

$$\begin{aligned}
11^A & (19,25), (24,26), (44,47), (57,231), (251,253), (252,254) \\
11^N & (34,42), (35,43), (36,44), (39,45), (40,46), (41,47), (71,75), (73,74), (76,78), (151,152)
\end{aligned}$$

Similarly, within $\mathbb{K}_{12}$ we determine via the invariants the following 75 candidate mutant cliques, all of which turn out to be genuine mutant cliques. These are depicted in the Appendix as Figures 9–18. We mention that our list of mutant cliques is in agreement with that of Stoimenow [28].

$$\begin{aligned}
12^A & (7,14), (13,15), (29,113), (36,694), (44,64), (45,65), (48,60), (59,63), (67,136), (91,111), \\
& (101,115), (102,107), (108,120), (114,117), (126,132), (131,133), (134,188), (154,162), \\
& (164,166), (167,692), (195,693), (639,680), (675,688), (811,817), (829,832), (830,831), \\
& (844,846), (30,33,157), (116,122,182) \\
12^N & (21,29), (22,30), (23,31), (26,32), (27,33), (28,34), (55,223), (58,222), (59,220), (63,225), \\
& (64,261), (67,229), (85,130), (86,131), (87,132), (88,133), (89,134), (90,135), (91,136), \\
& (92,137), (93,138), (98,125), (99,126), (122,127), (123,128), (124,129), (205,226), (206,227), \\
& (207,228), (208,212), (209,213), (210,214), (231,232), (252,262), (255,263), (256,264), \\
& (364,365), (421,422), (553,556), (670,681), (671,682), (691,692), (693,696), (56,57,221), \\
& (60,61,219), (62,66,224)
\end{aligned}$$




Figure 2: Mutant cliques within $\mathbb{K}_{11}^A$.




Figure 3: Mutant cliques within $\mathbb{K}_{11}^N$.

Of interest is that a single pair of knots $(12_{119}^N, 12_{120}^N)$ are distinguished by all three polynomials, but not by the (apparent) hyperbolic volume (to 10 places, 9.8759424404). This pair is depicted in Figure 4; they are related by the exchange of a 2 tangle with a $1/2$ tangle. For knots of greater crossing numbers, the same phenomenon appears more frequently.




Figure 4: 12_{119}^N and 12_{120}^N are *not* mutants, but have the same hyperbolic volume.

The maximum size of mutant candidate cliques, and presumably also of true mutant cliques, rises with crossing number. Amongst the knots of 11 to 14 crossings, the maximum candidate clique sizes are respectively 2, 3, 6 and 8.

With respect to $\mathbb{K}_{11}$ and $\mathbb{K}_{12}$, we also make the following observations.

- All elements of each (candidate) mutant clique are of the same braid index: 4 for all those in $\mathbb{K}_{11}$, and 4, 5 or 6 for those in $\mathbb{K}_{12}$. (There is no reason to suppose that mutation preserves braid index more generally.)
- None of the mutant cliques and mutant clique candidates are 2-bridge knots. This reflects the fact that 2-bridge links are known to be invariant under mutation [24, p48].

4 Where LG fails to distinguish prime knots

We now wish to use these new evaluations to identify the point at which LG first fails to distinguish pairs of distinct prime knot types. Recall from [4] that LG is known to be complete for all prime knots (including reflections) of at most 10 crossings. Of course LG , in common with many well-known quantum link invariants, fails to distinguish mutant links.⁴ To illustrate this, it

⁴A quantum link invariant based on a tensor product of representations of a quantum (super)algebra will fail to distinguish mutant links if the decomposition of the tensor product into irreducible submodules contains no multiplicities.

Some quantum link invariants for which the decomposition contains multiplicities distinguish at least *some* mutants. For instance each invariant associated with the $SU_q(N)$ representation with Young diagram $[2, 1]$ distinguishes the Kinoshita–Terasaka mutants [27]. An example closer to LG is the invariant associated with the 8-dimensional $U_q[gl(2|1)]$ adjoint representation [12].

is demonstrated in [9] that LG assigns the same polynomials to the Kinoshita–Terasaka mutant pair of 11-crossing prime knots. Note that there are no mutant pairs of prime knots of less than 11 crossings [24, p44].

So, we seek pairs of distinct *nonmutant* prime knots undistinguished by LG . Our starting point is the following fact. Recent work by Ishii and Kanenobu [20] has determined a class of nonmutant pairs of prime knots undistinguished by LG ; the smallest example of these is a pair of 14-crossing knots ($14_{6955}^A, 14_{7393}^A$) described by 16-crossing projections.⁵ Thus, we need only search $\mathbb{K}_{11}$ – $\mathbb{K}_{13}$ to identify where LG *first* fails to distinguish nonmutant prime knots.

We find that, modulo mutation, LG is indeed complete over $\mathbb{K}_{11}$. Moreover, all 11-crossing prime knots are chiral, so we know that LG is complete, modulo mutation, for all prime knots including reflections, of up to 11 crossings.

Within $\mathbb{K}_{12}$, we discover only the following nonmutant LG -pairs:

$$(12_{341}^A, 12_{627}^A), \quad (12_{17}^N, 12_{584}^N), \quad (12_{90}^N, 12_{416}^N) \quad \text{and} \quad (12_{135}^N, 12_{416}^N),$$

which we depict in Figure 5. More precisely, LG cannot distinguish the triple $(12_{90}^N, 12_{135}^N, 12_{416}^N)$, but 12_{90}^N and 12_{135}^N are mutants.

The respective polynomials are listed below. Note that the first polynomial is palindromic in q , as expected as 12_{341}^A and 12_{627}^A are achiral. (Also note that neither of these are 2-bridge knots.)

12_{341}^A 12_{627}^A	$(12\overline{q}^6 + 476\overline{q}^4 + 2474\overline{q}^2 + 4081 + 2474q^2 + 476q^4 + 12q^6)$ $- (\overline{p}^2 + p^2) (103\overline{q}^5 + 1147\overline{q}^3 + 3229\overline{q} + 3229q + 1147q^3 + 103q^5)$ $+ (\overline{p}^4 + p^4) (10\overline{q}^6 + 303\overline{q}^4 + 1580\overline{q}^2 + 2623 + 1580q^2 + 303q^4 + 10q^6)$ $- (\overline{p}^6 + p^6) (37\overline{q}^5 + 449\overline{q}^3 + 1325\overline{q} + 1325q + 449q^3 + 37q^5)$ $+ (\overline{p}^8 + p^8) (58\overline{q}^4 + 389\overline{q}^2 + 688 + 389q^2 + 58q^4)$ $- (\overline{p}^{10} + p^{10}) (51\overline{q}^3 + 206\overline{q} + 206q + 51q^3)$ $+ (\overline{p}^{12} + p^{12}) (27\overline{q}^2 + 62 + 27q^2)$ $- (\overline{p}^{14} + p^{14}) (8\overline{q} + 8q)$ $+ (\overline{p}^{16} + p^{16}) (1)$
12_{17}^N 12_{584}^N	$(12\overline{q}^4 + 188\overline{q}^2 + 647 + 774q^2 + 296q^4 + 22q^6)$ $- (\overline{p}^2 + p^2) (44\overline{q}^3 + 328\overline{q} + 685q + 492q^3 + 93q^5 + 2q^7)$ $+ (\overline{p}^4 + p^4) (67\overline{q}^2 + 313 + 427q^2 + 172q^4 + 13q^6)$ $- (\overline{p}^6 + p^6) (56\overline{q} + 179q + 149q^3 + 26q^5)$ $+ (\overline{p}^8 + p^8) (28 + 58q^2 + 22q^4)$ $- (\overline{p}^{10} + p^{10}) (8q + 8q^3)$ $+ (\overline{p}^{12} + p^{12}) (q^2)$
12_{90}^N 12_{135}^N 12_{416}^N	$(2\overline{q}^8 + 72\overline{q}^6 + 292\overline{q}^4 + 201 + 374\overline{q}^2 + 32q^2)$ $- (\overline{p}^2 + p^2) (18\overline{q}^7 + 168\overline{q}^5 + 348\overline{q}^3 + 277\overline{q} + 83q + 4q^3)$ $+ (\overline{p}^4 + p^4) (2\overline{q}^8 + 64\overline{q}^6 + 240\overline{q}^4 + 268\overline{q}^2 + 118 + 13q^2)$ $- (\overline{p}^6 + p^6) (14\overline{q}^7 + 118\overline{q}^5 + 199\overline{q}^3 + 115\overline{q} + 20q)$ $+ (\overline{p}^8 + p^8) (1\overline{q}^8 + 36\overline{q}^6 + 111\overline{q}^4 + 86\overline{q}^2 + 2)$ $- (\overline{p}^{10} + p^{10}) (5\overline{q}^7 + 40\overline{q}^5 + 50\overline{q}^3 + 15\overline{q})$ $+ (\overline{p}^{12} + p^{12}) (7\overline{q}^6 + 20\overline{q}^4 + 9\overline{q}^2)$ $- (\overline{p}^{14} + p^{14}) (4\overline{q}^5 + 4\overline{q}^3)$ $+ (\overline{p}^{16} + p^{16}) (\overline{q}^4)$

⁵In their notation, these knots are respectively $K[\beta; \pm 2, \mp 2]$, where $\beta = \sigma_2^2 \sigma_1^{-2}$.




Figure 5: The nonmutant LG -pairs within $\mathbb{K}_{12}^N$. A mutation relating 12_{90}^N and 12_{135}^N is clearly visible.

Thus, we learn that the lowest crossing number of a nonmutant pair of prime knots indistinguishable by LG is 12. We remark that there are no examples of LG -pairs of knots of *different* crossing numbers or different alternatingness amongst the currently LG -evaluated prime knots. We also confirm that LG is not unity for any of these but the unknot.

Note that none of the four nonmutant LG -pairs within $\mathbb{K}_{12}$ are distinguished by either the HOMFLYPT or Kauffman polynomials (they are distinguished by the hyperbolic volume, which demonstrates that they are not mutants). More generally, it remains an open question whether LG ever fails to distinguish a pair of links distinguished by *either* the HOMFLYPT or the Kauffman polynomial (that is, Problem 6.4 of [20]).

5 Where LG first fails to distinguish reflections

It is known that LG is not always able to distinguish the reflections of chiral links. Specifically, Ishii and Kanenobu [20] have constructed an infinite family of chiral prime knots whose chirality is undetected by LG . The smallest example they provide is a prime knot with a 28-crossing projection.⁶ KNOTSCAPE cannot reduce this projection to one of less than 28 crossings, but does reduce it to the elegant diagram of Figure 6.

We next wish to discover where LG *first* fails to distinguish the chirality of a chiral prime knot, and to that end we search within $\mathbb{K}_{11}$ – $\mathbb{K}_{16}$ as we know that LG distinguishes the chiralities of all chiral prime knots of up to 10 crossings [4]. Clearly, we require foreknowledge of *which* knots are chiral, and fortunately

⁶In their notation, this knot is $L[\sigma_2^{-2}\sigma_1^2; \frac{1}{2}, \frac{1}{4}, -\frac{1}{2}, -\frac{1}{4}]$.




Figure 6: A 28-crossing projection of a chiral prime knot whose chirality is undetected by LG .

KNOTSCAPE, together with [13], provides us with information of the symmetry classes (chiral or not, invertible or not) of the HTW knots.

Firstly, KNOTSCAPE uses the embedded SNAPPEA to determine the symmetry classes of the hyperbolic HTW knots. SNAPPEA determines the symmetry class of a given hyperbolic knot K via construction of its symmetry group $\text{Sym}(S^3, K)$, that is, the group of homotopy-equivalence classes of homeomorphisms of (S^3, K) . The symmetry group is unlikely to be discernible by casual inspection of an arbitrary projection; indeed, for any given link, it remains an open question as to whether there exists a (maximally-symmetric) projection in which the symmetry group is visible. It turns out that the symmetry groups of hyperbolic knots are all finite groups, either cyclic or dihedral [24, Chapter 10], as per those of the regular plane figures. Rather than by inspection of diagrams, SNAPPEA actually computes $\text{Sym}(S^3, K)$ from the hyperbolic decomposition of the knot complement $S^3 - K$. In contrast, the symmetry groups of nonhyperbolic knots are more general. Those of the nonhyperbolic HTW knots are deduced manually in [13]; they are all dihedral groups, some finite (in particular each torus knot has group D_1), and some infinite. The nonhyperbolic HTW knots are all chiral and invertible.

So, amongst the chiral knots within each set $\mathbb{K}_c^P$, we search for knots whose LG polynomials are palindromic, indicating that they fail to distinguish chirality. In this manner, we quickly discover that LG distinguishes the chirality of all chiral prime knots of up to 12 crossings. However, we do not have evaluations of LG for any complete set $\mathbb{K}_c^P$ beyond that point, and we uncover no failures amongst the knots within $\mathbb{K}_{13}$ (all of which are chiral) for which we have evaluations of LG .

Instead, we adopt a different approach to the problem. We begin by observing that mutation can change a chiral knot into an achiral knot; an illustration of such a pair of 16-crossing mutant projections appears in [23] (this example is originally due to Kanenobu [22]). We here confirm that the pair are 16-crossing prime knots: the chiral is 16_{259088}^A and the achiral is 16_{259984}^A . This example illustrates a more general situation.

Lemma 1. *Mutation can change the symmetry group (and thus the symmetry class) of a link.*

The first example of this phenomenon amongst the prime knots is the sole instance of it within $\mathbb{K}_{11}$, in the pair $(11_{39}^N, 11_{45}^N)$. Although both are chiral, the former is noninvertible (with symmetry group $\mathbb{Z}_2$) and the latter invertible (with symmetry group D_1).

Now, say that we have a mutant pair of links, one chiral, the other achiral. Any mutation-insensitive link invariant will necessarily assign the same value to the achiral link and to both the chiralities of the chiral link, and thus fail to distinguish the chirality of the latter. Thus, the fact that mutation can change the symmetry group of a link means that a link invariant which cannot distinguish mutants will also fail to distinguish some chiral links from their reflections. Incompleteness for mutation implies incompleteness for chirality. We formalise this as the following lemma.

Lemma 2. *Any mutation-insensitive link invariant is necessarily also incomplete for chirality.*

So, we seek an example of such a mutant pair of minimal crossing number amongst the prime knots. To that end, we may at first examine *candidate* mutant cliques (that is, cliques of common (apparent) hyperbolic volume, HOM-FLYPT and Kauffman polynomial), and we don't need to prove that a given chiral knot is a member of a mutant clique before we examine its LG -polynomial. Thus, we search the HTW knots for candidate mutant cliques containing at least one chiral and at least one achiral element, and of them, we need only the chiral elements. The first point at which we find such cliques is at 14 crossings. (There are no achiral 13-crossing prime knots.) Curiously, all such 14-crossing candidate mutant cliques manifest in triples, each containing exactly one chiral element. Inspection demonstrates that all candidates actually are mutants. Moreover, as oriented knots, all the elements of the triples are noninvertible, and so the achiral are actually all negative achiral, meaning that they are equal to their inverse reflections, but not equal to their reflections per se. We list the 13 triples found in Table 4; their diagrams are included in the draft of this paper lodged with the LANL arXiv.

So, the first point at which mutation changes a chiral prime knot into an achiral prime knot is at 14 crossings. Thus, we have the following.

Lemma 3. *Any mutation-insensitive link invariant will fail to distinguish the chirality of some 14-crossing chiral prime knots.*

Such an invariant will also fail for other prime knots, some of which may be of lower crossing number. The best that can be hoped for is that it is able to distinguish the chirality of all chiral prime knots of up to 13 crossings. We illustrate using the chiral 14_{13107}^A and the achiral 14_{13109}^A . A mutation relating them is visible in Figure 7. Indeed, LG cannot distinguish this pair, and is insensitive to the chirality of 14_{13107}^A , yielding the following palindromic polynomial.

chiral	achiral mutants	
14_{506}^A	14_{486}^A	14_{731}^A
14_{680}^A	14_{509}^A	14_{585}^A
14_{12813}^A	14_{12807}^A	14_{12875}^A
14_{12858}^A	14_{12815}^A	14_{12830}^A
14_{13107}^A	14_{13109}^A	14_{13489}^A
14_{13262}^A	14_{13269}^A	14_{13506}^A
14_{14042}^A	14_{14043}^A	14_{14671}^A
14_{17268}^A	14_{17265}^A	14_{17275}^A
14_{17533}^A	14_{17531}^A	14_{17680}^A
14_{1309}^N	14_{1327}^N	14_{1497}^N
14_{1641}^N	14_{1552}^N	14_{2132}^N
14_{1644}^N	14_{1555}^N	14_{1671}^N
14_{2164}^N	14_{1669}^N	14_{1925}^N

Table 4: Mutant cliques within $\mathbb{K}_{14}$ containing both chiral and achiral elements.

$$\begin{aligned}
& (348\overline{q}^6 + 5106\overline{q}^4 + 19778\overline{q}^2 + 30201 + 19778q^2 + 5106q^4 + 348q^6) \\
- & (\overline{p}^2 + p^2) (33\overline{q}^7 + 1525\overline{q}^5 + 10516\overline{q}^3 + 25021\overline{q} + 25021q + 10516q^3 + 1525q^5 + 33q^7) \\
+ & (\overline{p}^4 + p^4) (248\overline{q}^6 + 3615\overline{q}^4 + 14119\overline{q}^2 + 21641 + 14119q^2 + 3615q^4 + 248q^6) \\
- & (\overline{p}^6 + p^6) (15\overline{q}^7 + 724\overline{q}^5 + 5241\overline{q}^3 + 12779\overline{q} + 12779q + 5241q^3 + 724q^5 + 15q^7) \\
+ & (\overline{p}^8 + p^4) (62\overline{q}^6 + 1156\overline{q}^4 + 4982\overline{q}^2 + 7861 + 4982q^2 + 1156q^4 + 62q^6) \\
- & (\overline{p}^{10} + p^{10}) (113\overline{q}^5 + 1159\overline{q}^3 + 3182\overline{q} + 3182q + 1159q^3 + 113q^5) \\
+ & (\overline{p}^{12} + p^{12}) (121\overline{q}^4 + 766\overline{q}^2 + 1326 + 766q^2 + 121q^4) \\
- & (\overline{p}^{14} + p^{14}) (83\overline{q}^3 + 326\overline{q} + 326q + 83q^3) \\
+ & (\overline{p}^{16} + p^{16}) (36\overline{q}^2 + 81 + 36q^2) \\
- & (\overline{p}^{18} + p^{18}) (9\overline{q} + 9q) \\
+ & (\overline{p}^{20} + p^{20}) (1)
\end{aligned}$$




Figure 7: The chiral 14_{13107}^A and the achiral 14_{13109}^A are mutants.

So, as each $LG^{m,n}$ is insensitive to mutation, it will certainly fail to detect the chirality of some 14-crossing prime knots. In particular, we now have

14 as an upper bound on the point at which $LG^{2,1}$ first fails to distinguish the chirality of a chiral prime knot. Note that whilst the mutation-changing-chirality-to-achirality phenomenon doesn't occur within $\mathbb{K}_{13}$ (all the knots of which are chiral), there may still be a 13-crossing prime knot whose chirality is undistinguished by LG . Our evaluations of LG for 25% of these knots haven't uncovered such an example; indeed a wider search within $\mathbb{K}_{13}$ of the LG -feasible knots whose chirality is undetected by the HOMFLYPT or Kauffman polynomials also uncovers no such examples. Moreover, of the 53,418 HTW knots for which we have evaluations of LG (most of which are chiral), the *only* examples of chiral knots whose chirality is undetected by LG are 8 already contained within Table 4:

$$14_{680}^A, 14_{12813}^A, 14_{12858}^A, 14_{13107}^A, 14_{13262}^A, 14_{17268}^A, 14_{1309}^N, 14_{2164}^N.$$

In contrast to the statements about LG , the first example of a chiral prime knot whose chirality is undetected by both the HOMFLYPT and Kauffman polynomials is the well-known case $9_4^N \equiv 9_{42}$. (This is the only such case where *either* polynomial fails within $\mathbb{K}_9$.) We know that the chirality of 9_4^N is detected by LG , and that it has no mutants amongst the HTW prime knots. We then conclude with some open questions.

1. Does there exist a chiral prime knot whose chirality is undetected by LG yet is detected by the HOMFLYPT or Kauffman polynomials?
2. Can we find a chiral link whose chirality is undetected by LG yet is *not* a mutant of an achiral link? Recalling that we don't know whether mutation preserves crossing number or alternatingness, an easier version of this problem is to find a chiral prime knot whose chirality is undetected by LG and has no mutants of the same alternatingness and crossing number.
3. Recall that the question of whether there exists a chiral link which is mutation-equivalent to its reflection appears to remain unanswered. Can we find such a link? As every mutation-insensitive link invariant will be unable to detect the chirality of such a link, and as LG detects the chirality of all prime knots of up to 12 crossings, there can be no such links amongst the prime knots of up to 12 crossings. Candidates for such links might be found amongst the chiral links whose chirality is undistinguished by all mutation-insensitive link invariants (for example 14_{13107}^A). Then again, are such links *necessarily* mutants of their reflections?

Acknowledgements

The knot diagrams were made by KNOTSCAPE. Jon Links thanks the Australian Research Council for support through an Australian Research Fellowship. The authors thank Taizo Kanenobu for the keen observation of an error in an earlier draft of this paper.

References

- [1] Colin C Adams. *The Knot Book: an Elementary Introduction to Mathematical Theory of Knots*. Freeman, 1994.
- [2] David De Wit. Braids for odd pretzel knots. In preparation.
- [3] David De Wit. Walking through braided streams: Algorithms for the evaluation of state models for link invariants. In preparation.
- [4] David De Wit. Automatic evaluation of the Links–Gould invariant for all prime knots of up to 10 crossings. *Journal of Knot Theory and its Ramifications*, 9(3):311–339, May 2000. RIMS-1235, [math/9906059](#).
- [5] David De Wit. An infinite suite of Links–Gould invariants. *Journal of Knot Theory and its Ramifications*, 10(1):37–62, February 2001. [math/0004170](#).
- [6] David De Wit. Automatic construction of explicit R matrices for the one-parameter families of irreducible typical highest weight $(\dot{0}_m|\dot{\alpha}_n)$ representations of $U_q[gl(m|n)]$. *Computer Physics Communications*, 145(2):205–255, 15 May 2002. [math/0101027](#).
- [7] David De Wit. The 2-bridge knots of up to 16 crossings. Under consideration. [math/0409351](#), September 2004.
- [8] David De Wit, Atsushi Ishii, and Jon R Links. Infinitely many two-variable generalisations of the Alexander–Conway polynomial. Under consideration. [math/0405403](#), May 2004.
- [9] David De Wit, Louis H Kauffman, and Jon R Links. On the Links–Gould invariant of links. *Journal of Knot Theory and its Ramifications*, 8(2):165–199, March 1999. [math/9811128](#).
- [10] Thomas A Gittings. Minimum braids: A complete invariant of knots and links. [math/0401051](#), 2004.
- [11] Cameron McA Gordon and John Luecke. Knots are determined by their complements. *Journal of the American Mathematical Society*, 2:371–415, 1989.
- [12] Mark D Gould, Jon R Links, and Yao-Zhong Zhang. Type-I quantum superalgebras, q -supertrace and two-variable link polynomials. *Journal of Mathematical Physics*, 37:987–1003, 1996.
- [13] Jim Hoste, Morwen Thistlethwaite, and Jeff Weeks. The first 1,701,936 knots. *The Mathematical Intelligencer*, 20(4):33–48, 1998.
- [14] Noriko Imafuji and Mitsuyuki Ochiai. Computer aided knot theory using Mathematica and Mathlink. *Journal of Knot Theory and its Ramifications*, 11(6):945–954, September 2002.

- [15] Atsushi Ishii. The Links–Gould invariants of the Kanenobu knots. *Kobe Journal of Mathematics*, 20:53–61, 2003.
- [16] Atsushi Ishii. Algebraic links and skein relations of the Links–Gould invariant. *Proceedings of the American Mathematical Society*, 132:3741–3749, 2004.
- [17] Atsushi Ishii. The LG polynomial as a generalization of the Alexander–Conway polynomial. Under consideration, 2004.
- [18] Atsushi Ishii. The Links–Gould invariant as Vassiliev invariants. Under consideration, 2004.
- [19] Atsushi Ishii. The Links–Gould invariant of closed 3-braids. *Journal of Knot Theory and its Ramifications*, 13(1):41–56, February 2004.
- [20] Atsushi Ishii and Taizo Kanenobu. Different links with the same Links–Gould invariant. To appear in *Osaka Journal of Mathematics*, 2004.
- [21] Atsushi Ishii and Taizo Kanenobu. A relation between the Links–Gould invariant and the Kauffman polynomial. Under consideration, 2004.
- [22] Taizo Kanenobu. A note on 2-fold branched covering spaces of S^3 . *Mathematische Annalen*, 256:449–452, 1981.
- [23] Louis H Kauffman. An invariant of regular isotopy. *Transactions of the American Mathematical Society*, 318(2):417–471, April 1990.
- [24] Akio Kawauchi. *A Survey of Knot Theory*. Birkhäuser Verlag, 1996.
- [25] Jon R Links and Mark D Gould. Two variable link polynomials from quantum supergroups. *Letters in Mathematical Physics*, 26(3):187–198, November 1992.
- [26] Charles Livingston. TABLE OF KNOT INVARIANTS.
<www.indiana.edu/~knotinfo/>.
- [27] Hugh R Morton and Peter R Cromwell. Distinguishing mutants by knot polynomials. *Journal of Knot Theory and its Ramifications*, 5(2):225–238, 1996.
- [28] Alexander Stoimenow. KNOT DATA TABLES.
<www.math.toronto.edu/stoimeno/ptab/index.html>.
- [29] Pierre Vogel. Representation of links by braids: a new algorithm. *Commentarii Mathematici Helvetici*, 65(1):104–113, 1990.
- [30] Stephan Wehrli. Khovanov homology and Conway mutation.
math/0301312, 2003.

Appendix

On the following pages are illustrated some mutant cliques mentioned within the main text. Figures 9–18 illustrate the mutant cliques within $\mathbb{K}_{12}$, and Figures 19–21 illustrate the mutant cliques containing both chiral and achiral knots within $\mathbb{K}_{14}$.




Figure 8: The example of Kanenobu of a chiral-achiral pair of mutant prime knots: the chiral 16^A_{259088} and the achiral 16^A_{259984} .




Figure 9: Alternating 12-crossing mutant cliques 1/4




Figure 10: Alternating 12-crossing mutant cliques 2/4




Figure 11: Alternating 12-crossing mutant cliques 3/4




Figure 12: Alternating 12-crossing mutant cliques 4/4




Figure 13: Nonalternating 12-crossing mutant cliques 1/6




Figure 14: Nonalternating 12-crossing mutant cliques 2/6




Figure 15: Nonalternating 12-crossing mutant cliques 3/6




Figure 16: Nonalternating 12-crossing mutant cliques 4/6




Figure 17: Nonalternating 12-crossing mutant cliques 5/6




Figure 18: Nonalternating 12-crossing mutant cliques 6/6




Figure 19: Alternating 14-crossing mutant cliques containing both chiral and achiral elements $1/2$.




Figure 20: Alternating 14-crossing mutant cliques containing both chiral and achiral elements $2/2$.




Figure 21: Nonalternating 14-crossing mutant cliques containing both chiral and achiral elements.